

COVID Vaccine Injury Reports Grow in Number, But Trends Remain Consistent

Data released today by the CDC confirm several ongoing trends, including that 47% of deaths occurred in people who reported becoming sick within 48 hours of receiving a COVID vaccine, and 20% of deaths were cardiac-related.

By Megan Redshaw, March 5, 2021

According to data released today by the Centers for Disease Control and Prevention (CDC), the number of injuries and deaths reported to the [Vaccine Adverse Event Reporting System \(VAERS\)](#) following COVID vaccines continues to climb.

Between Dec. 14, 2020, and Feb. 26, a total of [25,212 total adverse events](#) were reported to VAERS, including [1,265 deaths](#) and [4,424 serious injuries](#).

From the [2/26/2021 release of VAERS data](#):

Found 25,212 cases where Vaccine is COVID19

Table

↓	↑ ↓	
Event Outcome	Count	Percent
Death	1,265	5.02%
Permanent Disability	479	1.9%
Office Visit	3,888	15.42%
Emergency Room	22	0.09%
Emergency Doctor/Room	4,908	19.47%
Hospitalized	2,742	10.88%
Hospitalized, Prolonged	1	0%
Recovered	9,236	36.63%
Birth Defect	36	0.14%
Life Threatening	886	3.51%
Not Serious	8,946	35.48%
TOTAL	† 32,409	† 128.55%

† Because some cases have multiple vaccinations and symptoms, a single case can account for multiple entries in this table. This is the reason why the Total Count is greater than 25212 (the number of cases found), and the Total Percentage is greater than 100.

In the U.S., [70.45 million](#) COVID vaccine doses had been administered as of Feb. 26.

[VAERS](#) is the primary mechanism for reporting adverse vaccine reactions in the U.S. Reports submitted to VAERS require further investigation before confirmation can be made that the reported adverse event was caused by the vaccine.

According to the latest data, [1,136 of 1,265 reported deaths](#) were in the U.S. Of the total, 31% of the deaths occurred within 48 hours of vaccination, and 47% of deaths occurred in people who became ill within 48 hours of being vaccinated. Twenty percent of deaths were related to cardiac disorder.

Fifty-three percent of those who died were male, 45% were female and the remaining death reports did not include gender of the deceased. The [average age](#) of those who died was 77.8 and the youngest death confirmed was a 23-year-old.

As of Feb. 26, [180 pregnant women](#) had reported adverse reactions to COVID vaccines, including 56 reports of [miscarriage or premature birth](#). None of the COVID vaccines approved for [Emergency Use Authorization](#) (EUA) have been tested for safety or efficacy in pregnant women. Yet health officials are urging pregnant women to get the vaccine, and many are enthusiastically doing so. As The Defender reported:

“Even without data from [Pfizer](#) or [Moderna](#) sufficient to inform vaccine-associated risks in pregnancy, [expectant](#) doctors, nurses and others appear eager for the shots, perhaps influenced by the American College of Obstetricians and Gynecologists, [which states](#) that ‘neither a conversation with a clinician nor even a pregnancy test are necessary prerequisites.’”

[Continue reading here...](#)